No.



830011

HHE UNIVIED SHAVES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Nickerson American Plant Breeders, Inc.

Withereas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE; IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF EIGHTERN YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT TY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS Y THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT

'Ram'

In Testimony Wathereof, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington this 30th day of August in the year of our Lord one thousand nine hundred and eighty-five.

Secretary of Agriculture

Start.

Senth N. Evan Commissioner Plant Variety Protection Office

Plant Variety Protection Office Agricultural Marketina Service

	AGRICULTURAL MAR	RETING SERVICE			FORM APPROVE OMB NO. 40-R382
	APPLICATION FOR PLANT VARI		N CERTIFICATE		ent variety protection ma empleted application for U.S.C. 553).
	1a. TEMPORARY DESIGNATION OF VARIETY	16. VARIETY NAME	· · · · · · · · · · · · · · · · · · ·	FOR OFFICE	IAL USE ONLY
•	HW77-391S4	Ram		PV NUMBER 830	0111
	2. KIND NAME	3. GENUS AND SPEC	CIES NAME	FILING DATE	TIME XXX
	Hard Red Winter Wheat	Triticum ae	stivum	4/11/83	3:00 P.M
	4. FAMILY NAME (BOTANICAL)	5. DATE OF DETER		FEE RECEIVED	DATE /. /11 /00
	Gramineae	1) September 2) September	1979	\$\frac{1,000}{500.00}	4/11/83 7/16/85
Eigh	6. NAME OF APPLICANT(S) NICKERS& D	7. ADDRESS (Street Code)	and No. or R.F.D. No.,	City, State, and ZIP	B. TELEPHONE AREA
131 8 8 R	North American Plant Breeders,	, ,	hnson Dr., P.0 , KS 66201	. Box 2955	913-384-4940 KS 303-532-3721 CC
3.61	9. IF THE NAMED APPLICANT IS NOT A PORGANIZATION: (Corporation, partners	PERSON, FORM OF hip, association, etc.)	10. IF INCORPORAT DATE OF INCOR	ED, GIVE STATE AND PORATION	11. DATE OF INCOR- PORATION
e .	Corporation		Delaware, 1-		1-19-83
ZE S	12. NAME AND MAILING ADDRESS OF APP	LICANT REPRESENTA	TIVE ST. FARRING	or C. Bruns	ATION AND RECEIVE
3 4	P.O. Box 2955		P.O. Box 30	••• •	
A. S.	*	66201	Berthoud, C	0 80513	
	13. CHECK BOX BELOW FOR EACH ATTAC				-
	X 13A. Exhibit A, Origin and Bre	eding History of the V	ariety (See Section 5	52 of the Plant Variety	Protection Act.)
	13B. Exhibit B, Novelty States	nent.	•		
•	X 13C. Exhibit C, Objective Desc	ription of the Variety	(Request form from	Plant Variety Protecti	ion Office.)
	X 13D. Exhibit D, Additional Des X 13E. Exhibit E, Quali	ty Data and Sta	tistical Data.		
	148. DOES THE APPLICANT(S) SPECIFY THA SEED7- (See Section 83(a). (If "Yes," ansu	T SEED OF THIS VARIE ver 14B and 14C below.)		RIETY NAME ONLY AS NO	A CLASS OF CERTIFIE
	14b. DOES THE APPLICANT(S) SPECIFY THA		TION BEYOND B		
	X YES NO		[X] FOUNDATION	REGISTERED	X CERTIFIED
	15a. DID THE APPLICANT(S) FILE FOR PROT name of countries and dates.)	ECTION OF THIS VARI	ETY IN OTHER COU	NTRIES? TYES	NO (If "Yes," give
	·				
	15b. HAVE RIGHTS BEEN GRANTED THIS V. and dates.)	ARIETY IN OTHER COL	INTRIES? TYES	NO (If "Yes,"	give name of countries
	and dates, y				
•	16. DOES THE APPLICANT(S) AGREE TO TH	E PUBLICATION OF HI	S/HER (THEIR) NAME	(S) AND ADDRESS IN	THE OFFICIAL
	17. The applicant(s) declare(s) that a viable replenished upon request in accordance	e sample of basic seed e with such regulation	of this variety will be s as may be applicabl	e furnished with the a	pplication and will be
	The undersigned applicant(s) is (are) the variety is distinct, uniform, and stable	he owner(s) of this sex	ually reproduced nov	vel plant variety, and l	believe(s) that the
	42 of the Plant Variety Act.		•		
	Applicant(s) is (are) informed that fals	e representation hereir	i can jeopardize prot	ection and result in po	enalties.
•.	(DATE)	•	Cours	IGNÁTURE OF APPLIC	CANT)
•	March 28, 1983		Olh	son	
	(DATE) ' FORM GR-470 (1-78)		Tyl is	GNATURE OF APPLIC	CANT) 1
	•		RI	ECEIVED MAR	2 8 1923

Exhibit A

Origin and Breeding History of Ram

PEDIGREE: II18889/Trapper//C0652643/3/Baca

DATE OF CROSS: 1973

HISTORY: The breeding history of Ram started in 1973 with the cross of CO701411 (F6) and Baca. This F1 was increased in 1974, and grown as an F2 population in 1975. Single rows of F3 lines were grown in 1976 at three locations. One of these lines was advanced into regional yield trials in 1977. In 1979, head-rows were grown at Berthoud, Colorado. One of these rows was distinctly taller than the majority and was harvested separately and given the designation HW77-391S4. Ram was put into

regional yield testing in 1981 and breeders seed increases were initiated. Approximately 1370 units of foundation seed were produced in 1982.

Ram is uniform and stable. Less than 1% of the plants have been rogued from registered fields in 1982. Approximately 90% of these rogued plants have been three to fourteen centimeters taller than Ram. Less than 1% of these taller plants may be encountered in subsequent generations.

Exhibit B

Novelty Statement

Ram is most similar to the hard red winter wheat, Hawk. However, it can be distinguished on the following morphological characteristics:

- Ram and Hawk differ significantly in glume width; Ram's are narrower (see supporting statistical data, Exhibit E).
- Ram and Hawk differ significantly in height; Ram is taller (see supporting statistical data, Exhibit E).
- Ram has a brown phenol reaction, color category #4. Hawk displays 80% light brown (#3) phenol reaction and 20% brown (#4) phenol reaction.

Anova Table for Height Ram versus Hawk

Source	df	ss	ms
Total	39	9629.5	
Rep	19	8875.5	467.1
VAR	İ	562.5	562.5**
Error	19	191.5	10.1

F test (.05) = 55.7**

** There is a significant difference at the 5% and 1% levels.

variety	mean (cm)
Hawk	75.5
Ram	83.0

Anova Table for Glume Width Ram versus Hawk

Source	df	SS	ms
Total	49	2.960	·
VAR	7	1.312	1.312**
Error (Res)	48	1.648	.034

F test = 38.59**

** The difference in glume width between Hawk and Ram is significant at the 5% and 1% levels.

variety	•	mean (mm)
Hawk		3.45
Ram		3.12

FORM GR-470-6 (2-15-73)

UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN DIVISION

EXHIBIT C (Wheat)

HYATTSVILLE, MARYLAND 20782

OBJECTIVE DESCRIPTION OF VARIETY

WHEAT (TRITICUM SPP.) INSTRUCTIONS: See Reverse. NATEREASSING ANTIST North American Plant Breeders, INC. FOR OFFICIAL USE ONLY 8300111 ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code) VARIETY NAME OR TEMPORARY DESIGNATION 5201 Johnson Dr., P.O. Box 2955 Mission, KS 66201 RAM Place the appropriate number that describes the varietal character of this variety in the boxes below. Place a zero in first box (e-g- 0 8 9 or 0 9) when number is either 99 or less or 9 or less. I, KIND: I = COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5 = POLISH 6 = POULARD 7 ≈ CLUB 2. TYPE: 3 = OTHER (Specify) l = SOFT2 1 = SPRING 2 = WINTER 3 = OTHER (Specify) 2 = HARD 2 = RED 3 = OTHER (Specify) 3. SEASON - NUMBER OF DAYS FROM TO: planting FIRST FLOWERING LAST FLOWERING MATURITY (50% Flowering): NO. OF DAYS EARLIER THAN] = ARTHUR 2 = SCOUT3 = CHRIS4 = LEMHI 5 = NUGAINES 6 = LEEDS 0 5. PLANT HEIGHT (From soil level to top of head): 0 CM. HIGH 2 = SCOUT3 = CHRIS CM. TALLER THAN ... 1 = ARTHUR 6 = LEEDS 4 = LEMHI5 = NUGAINES CM. SHORTER THAN 6. PLANT COLOR AT BOOTING (See reverse): 7. ANTHER COLOR:

1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN

I = YELLOW 2 = PURPLE

8. STEM:

Anthocyanin: I = ABSENT

2 = PRESENT

2

Waxy bloom: } = ABSENT

2 = PRESENT

Hairiness of last

internode of rachis: 1 = ABSENT

2 = PRESENT

Internodes: 1 = HOLLOW

NO. OF NODES (Originating from node above ground)

CM, INTERNODE LENGTH BETWEEN FLAG LEAF AND LEAF BELOW

9. AURICLES:

Anthocyanin: 1 = ABSENT

2 = PRESENT

Hairiness: 1 = ABSENT

2 = PRESENT

10. LEAF:

booting stage: -

] = ERECT 2 = RECURVED 3 = OTHER (Specify):___

Flag leaf: 1 = NOT TWISTED 2 = TWISTED

2 = PRESENT

Hairs of first leaf sheath: 1 = ABSENT

2 = PRESENT

Waxy bloom of flag leaf sheath: 1 = ABSENT

MM. LEAF WIDTH (First leaf below flag leaf)

CM. LEAF LENGTH (First leaf below flag leaf):

Ram			8300111
FORM GR-470-6 (REVERS	,E)		
3 Density: 1 = LAX	2 = DENSE 3=middense ave. 40.4	Shape: = TAPER 4 = OTHER	RING 2 = STRAP 3 = CLAVATE R (Specify)
4 Awnedness: 1 = Av		3 = AWNLETED 4 = AWNE	ם:
Color at maturity: 5	I=white 2=YELLOW 3=PINK 4 5=brown 6=black 7=oth	= RED ER (Specify):	<u>.</u>
6 6 CM. LENGTH		1 1 mm. WIDTH	· · · · · · · · · · · · · · · · · · ·
12. GLUMES AT MATUR 2 Length: 1 = SHORT 3 = LONG	_		w(CA. 3 mm.) 2 = MEDIUM(CA. 3.5 mm.) (A. 4 mm.) (average 6.78 mm
1 .3 [ring 2 = oblique 3 = rounded are 5 = elevated 6 = apiculate	Beak: 1 = OBTUSE),
13. COLEOPTILE COLOR	₹:	14. SEEDLING ANTHOCY	ANIN:
1 = WHITE 2 = R	RED 3 = PURPLE	2 I = ABSENT	epresent (slight)
15. JUVENILE PLANT GR	ROWTH HABIT:		
2 I = PROSTRATE	2 = SEMI-ERECT. 3 = EREC	ST.	
16. SEED:			
3 Shape: 1 = OVATE	2 = OVAL 3 = ELLIPTICAL	Cheek: 1 = ROUND	ED 2 = ANGULAR
2 Brush: 1 = SHORT	midlong 2 = 3 = Long	Brush: I = NOT C	DLLARED 2 = COLLARED
Phenol reaction (See instructions):	1 = IVORY . 2 = FAWN 3 = LT. BROW 4 = BROWN 5 = BLACK	N	
3 Color: 1 = WHITE	2 = AMBER 3 = RED 4 = PURPLE	5 = OTHER (Specify)	
6 6 MM. LENGTH	3 3 MM. WIDTH	4 5 GM. PER 1000	SEEDS
17. SEED CREASE:		• .	
· •	LESS OF KERNEL 'WINOKA'	, , , ,	R LESS OF KERNEL 'SCOUT'
	ESS OF KERNEL 'CHRIS' AS WIDE AS KERNEL 'LEMHI'	= :-	LESS OF KERNEL 'CHRIS' LESS OF KERNEL 'LEMHI'
		moderate susceptib	
4 (Races) field mi	1 FAF BUST	O STRIPE RUST (Races)	1 LOOSE SMUT
ture of 15TNM	1.	· 	il borne mosaic virus
	ed, 1 = Susceptible, 2 = Resistant) 3)		
0 SAWFLY	APHID (Bydv.)	GREEN BUG	O CEREAL LEAF BEETLE
OTHER (Specify)	HESSIAN FLY		
	RACES:] D] E] F] G
O. INDICATE WHICH VARI	ETY MOST CLOSELY RESEMBLES THAT S	HEMITTED.	
CHARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Plant tillering	Hawk	Seed size	Hawk
Leaf size	Hawk	Seed shape	Hawk
Leaf color	Hawk	Coleoptile elongation	Hawk
Leaf carriage	Hawk	Seedling pigmentation	Hawk -

INSTRUCTIONS

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

Exhibit D

Additional Description

Ram is a hard red winter wheat tested as HW77-391S4. It was developed by North American Plant Breeders.

Ram is an intermediate to tall height semidwarf variety with good straw strength, intermediate to late maturity and good winterhardiness. Milling and baking properties are acceptable.

Juvenile plant growth habit is semi-erect. Plant color at boot is green with an erect, twisted flag leaf. Head shape is tapering to strap, middense, awned and head color is white at maturity. Glumes are of medium length and narrow width with rounded to square shoulders and acuminate beaks; seed shape is elliptical. Seed crease is narrow and depth is shallow.

Ram is adapted to the panhandles of Oklahoma, Texas and Nebraska, western Kansas and eastern Colorado.

SAMPLE TEST WHT FLR FLR NAME NAME T. PROT YLD PROT T. PROT YLD PROT T. PROT YLD PROT T. PROT T	(North A	imer ican	North American Plant Breeders	kreeder	'n							
SAMPLE LOC WIT. PROIT YLD PROIT ASH CURNE NAME LOS WIT. PROIT YLD PROIT ASH CURNE NAME LOC WIT. PROIT YLD PROIT ASH CURNE NAME LOS WIT. PROIT YLD PROIT ASH CURNE NAME LOC WIT. PROIT YLD PROIT ASH CURNE NILL ASH CURNE NAME LOC WIT. PROIT YLD PROIT ASH CURNE NILL ASH	8			٠.			HARD RE	E WINTE	IR WHEAT	GUALI	Ł						PAG	
THE LOC WIT, PROT TEST WHT FLR FLR FLR FLR FLR MIX MIX DOUGH LORF GRING TEXT CROWN TEXT COLOR SCORE SC			1	MHEA	TFL0	UR QUA	LITY	į			SAKING	ผมคนา	} .					
1b/8u 14xmb x 14xmb R x min R cc R R R R R 773-C 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	SAMPLE	LOC	TEST WT.	PROT	FLR YLD	FLR PROT	FLR ASH	MIX	ABS.	1	1	VOP T				MILL	RAKE	TOTAL
7-39154 CK 53.3 11.7 71.9 10.4 0.393 6 59.0 6.0 8 980 6 7 9 72-C 73-C 11.7 71.9 10.4 0.393 6 59.0 6.0 4.5 8 980 6 7 9 775 6 73-C 11.5 7-39154 HN 58.1 12.5 70.6 11.6 0.474 6 59.0 4.5 8 870 9 8 7 9 7 9 69-D 77-C 1 75-C 1 75-C 1 1.2 72.1 10.7 0.404 4 57.0 3.5 8 810 8 6 7 9 9 7 7 69-D 77-C 1 75-C 1 75-C 1 1.2 72.1 10.7 0.404 4 57.0 3.5 8 810 8 8 7 9 7 7 89-D 77-C 1 1 7-39154 HN 58.1 12.3 70.9 11.3 0.406 6 61.0 5.5 9 863 8 7 8 7 8 7 7 8 863 8 7 7 8 70-C 7 76-C 1 1 7-29154 HN 59.9 11.3 0.456 8 62.0 3.0 3.5 9 750 8 9 9 75-C 90-B 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		1	16/Bu	14%mb		14%mb	14%mb	œ		min	æ	2	æ	 ac	4		! ! !	1
7-39154 S0 58.6 12.0 67.8 11.5 0.434 6 60.0 4.5 8 775 8 8 775 8 8 775 7 7 7 7 7 7 7 7	W77-33154	ð	59,3	11.7	71.3	10.4	0.393	8	0.68	0.3	o,	000	ú		0	0 0	0	1 0
7-39154 LK 53.7 12.8 70.6 11.6 0,474 6 59.0 4.5 8 890 8 7 9 69-0 77-0 1 7-39154 HN 58.1 12.5 70.8 11.3 0,377 5 58.0 4.0 8 870 9 8 7 9 69-0 77-0 1 7-39154 HN 58.1 12.5 70.8 11.3 0,377 5 58.0 4.0 8 870 9 8 7 9 73-0 77-0 1 7-39154 CK 51.1 13.9 69.6 12.3 0,000 6 61.0 5.5 9 863 8 7 8 70-0 76-0 1 68-0 1 1 7-39154 CK 51.1 13.9 69.6 12.3 0,000 6 61.0 5.5 9 863 8 7 8 70-0 76-0 1 7-39154 CK 59.8 12.3 70.9 11.3 0,456 8 62.0 5.0 8 895 7 7 8 70-0 76-0 1 7-39154 CK 59.8 11.5 70.4 10.5 0,339 6 59.0 3.0 9 70-0 7 8 9 9 77-0 1 7-30-0 1 7-39154 CK 56.2 11.5 70.4 10.5 0,386 5 69.0 2.0 8 810 7 8 9 9 77-0 1 7-39154 CK 56.2 11.5 70.4 11.2 0,405 6 61.2 3.6 9 875 8 9 9 76-0 83-0 1 7-3 11.8 11.8 11.8 11.8 11.8 11.8 11.8 11.	W77-39154	90	58.6	12.0	£7.8	11.5	0.434	Ψ	60.0	1	joj	7.	5 d	. 0	. 0	ن ر ۱۰۱ ۱۰۷) () (
7-39154 HN 58.1 12.5 70.8 11.3 0.377 5 58.0 4.0 8 870 9 8 8 73-C 77-C 17-C 17-C 17-C 17-C 17-C 17-C 17	ヤビアーののエS4		53,7	12,8	70.8	11.6	0.474	Ø	90.00	4) o	000) e	A (o	100) (1 1 1) (0 1
7-39154 BR 53-3 11.2 72.1 10.7 0.404 4 57.0 3.5 9 910 8 6 7 69-D 69-D 7-39154 CK 51.1 13.9 69-6 12.3 0.000 6 61.0 5.5 9 850 7 8 9 71-C 78-C 1 AVERAGE 56.8 12.4 70.5 11.3 0.347 6 59.0 4.7 8 863 8 7 8 70-C 76-C 76-C 76-C 76-C 76-C 76-C 76-C 76	W77-39154		58.1	12.5	20.8	11.3	0.377	ហ	58.0	0.4	, tō3	0.70	o	. o;	ìα	ن ن م را م (د) (1 1	
AVERAGE CK 51.1 13.9 69.6 12.3 0.000 6 61.0 5.5 9 850 7 8 9 71-C 78-C 1 AVERAGE 56.8 12.4 70.5 11.3 0.347 6 59.0 4.7 8 863 8 7 8 70-C 76-C 76-C 76-C 10.0 11.3 0.347 6 59.0 4.7 8 863 8 7 8 70-C 76-C 76-C 10.0 11.3 0.456 8 62.0 3.5 9 75-C 8 8 9 77-C 10.0 10.0 10.0 10.0 10.0 10.0 10.0 10.	4M77-39154	噐	0.00 0.00	11.2	72.1	10.7	0.404	₫	57.0	ю 10	ŋ	810	· 00	o vo) K) ii	0-89	137-10
AVERAGE 56.8 12.4 70.5 11.3 0.347 6 59.0 4.7 8 863 8 7 8 70-C 76-C 76-C 76-C 76-C 76-C 76-C 76-C 76	HW77-39154		51.1	13.9	9.69	12.3	0.000	ம	61.0	ຊຸນ	ø	850	٨	\$	ሙ	71-C	78-C	149-C
TON CK 59-8 12.3 70.9 11.3 0.456 8 62.0 5.0 8 895 7 7 9 81-B 80-B 1 10	AVERAG	ш	56.8	12,4	70.5	11.3	0.347	Ø	23.0	7.	ø	863	ഇ	~1	ಖ	70-C	76-0	146-0
TON CK 59-8 12.3 70.9 11.3 0.456 8 62.0 5.0 8 895 7 7 9 81-B 80-B 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1										1		1			-			
TON LK 56.2 11.5 70.4 12.0 0.386 5 63.0 3.5 3 750 8 8 9 75-C 80-R 1 TON LK 59.4 11.5 70.4 12.0 0.386 5 63.0 3.5 3 80.5 3 9 72-C 78-C 1 TON ER 62.0 12.5 73.4 12.0 0.386 5 63.0 2.0 8 810 7 8 9 77-C 1 TON CK 56.2 11.5 70.3 10.1 0.397 5 60.0 4.3 9 830 8 9 63-D 81-B 1 AVERAGE 59.4 12.2 70.6 11.2 0.405 6 61.2 3.6 9 875 8 8 9 76-C 33-B A-EXCELLENT 8-GOOD C-ACCEPTABLE D-QUESTIONABLE F-UNACCEPTABLE	NOTMEN	ď	5.6 6.6	12.3	70 B	11,3	0.456	ø	62.0	5.0		8,95	٨	~	თ	81-B	80-B	161-B
TCN HN 59:9 11:5 73.4 10:5 0.339 6 59:0 3.5 8 975 8 8 9 72-C 78-C 100 HN 59:9 11:5 73.4 12:0 0.386 5 63:0 2.0 8 810 7 8 9 9 72-C 78-C 100 N BR 62:0 12:5 73.4 12:0 0.386 5 63:0 2.0 8 810 7 8 9 9 77-C 1 70 N CK 56:2 11:5 70:3 10:1 0.397 5 60:0 4.3 9 830 8 9 9 63-D 81-B 1 AVERAGE 59:4 12:2 70:6 11:2 0.405 6 61:2 3:6 9 875 8 9 76-C 83-B 1 A-EXCELLENT 8-600D C-ACCEPTABLE D-QUESTIONABLE F-UNACCEPTABLE		χ.	υ η 10 ο	6.0	က (ထို (1.0	0.400	i~ i	61.0	ო ი		230	ø	o)	o,	72-C	80-F	155-C
TON BR 62.0 12.5 73.4 12.0 0.339 6 59.0 3.5 8 875 8 8 9 72-C 78-C 1 70.8 11.5 70.3 10.1 0.397 5 60.0 4.3 9 830 8 9 9 63-D 81-B 1 AVERAGE 5 61.2 3.6 9 875 8 9 9 76-C 83-B 1 40-EXCELLENT 8-GOOD C-ACCEPTABLE D-QUESTIONABLE F-UNACCEPTABLE		5 :	9 (0	0.1	n .	27	0.44		62.0	о 0		+0001	ø	σι	Ф	79-C	90-A	168-8
TON CK 56.2 11.5 73.4 12.0 0.386 5 63.0 2.0 8 810 7 8 9 97-C 1 TON CK 56.2 11.5 70.3 10.1 0.397 5 60.0 4.3 9 830 8 9 9 63-D 81-B 1 AVERAGE 59.4 12.2 70.6 11.2 0.405 6 61.2 3.6 9 875 8 8 76-C 83-B A-EXCELLENT 8-GOOD C-ACCEPTABLE D-QUESTIONABLE F-UNACCEPTABLE		Z (n (0.11	4.0	10.5	0.333	ဖ	23.0	in in	₩	875	ø	σj	ሙ	, 72-C	7007	150-C
TON CK 56.2 11.5 70.3 10.1 0.397 5 60.0 4.3 9 830 8 9 9 63-D 81-B 1 AVERAGE 59.4 12.2 70.6 11.2 0.405 6 61.2 3.6 9 875 8 9 76-C 83-B A-EXCELLENT 8-GOOD C-ACCEPTABLE D-QUESTIONABLE F-UNACCEPTABLE	20 - MUZ	£	0.79	12.3	43.4	12.0	986.0	ហ	63.0	2.0	øj	810	ı.	ø	ტ	83-8	77-C	L60-B
AVERAGE 59.4 12.2 70.6 11.2 0.405 6 61.2 3.6 9 875 8 9 76-C 83-B A-EXCELLENT 8-GOOD C-ACCEPTABLE D-QUESTIONABLE F-UNACCEPTABLE	NEWTON	ኝ	56.2	11.5	70.3	10.1	0.397	n	60.09	4.9	m	830	ø	σı	ø	93-E	81-B	144-C
A-EXCELLENT B-GOOD C-ACCEPTABLE D-QUESTIONABLE	AVERAG	ш	4.60	12.2	70.6	11.2	0.405	O	61.2	9	ø,	87.5	σ	a)	ø	76-C	83-B	159-C
	GRADES: A	-באכברו	ENT	B-6000		ACCEPT	ABLE	ם-פר	ESTIONAL	U U	J-7	NACCES	TABLE					